

PAC
12/17/2019



For faster service, request your refills 2-3 days in advance, using any of these methods:

- Go to www.fruthpharmacy.com and press the button for prescription refills
- Call (304) 720-8341 and follow the prompts to use our automated refill line

PACHAURI, VEDANSHI

720 OXFORD CIR
CHARLESTON, WV 25314-1786
(304) 382-2939
DOB: 08/20/2012

RX# 6819365
CEFDINIR 250 MG/5ML SUS LUPI

GENERIC DISPENSED
NDC# 68180-0723-20 QTY 120
PRESCRIBER: RAHEEL KHAN
DAYS 10.00
REFILLS: 0 UNTIL 12/16/20
DISPENSED BY: SA
TX 1765178



AMOUNT DUE \$79.39

INS #1:UHCW

Fruth Prescription Rewards
NO CUSTOMER LOYALTY REWARDS NUMBER ON FILE FOR THIS PATIENT.
CASHIER: THIS PRESCRIPTION IS ELIGIBLE FOR REWARDS, PLEASE ASK FOR THE PATIENTS LOYALTY REWARDS CARD NUMBER AND UPDATE THE SYSTEM.

Tell us how we're doing, take our customer service survey at: <http://svy.mk/1LcUVPU> **25**

Pachauri, Vedanshi
CEFDINIR 250 MG/5ML SUS LUPI

GENERIC NAME: Cefdinir Suspension (SEF di ner)

COMMON USES: It is used to treat bacterial infections.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to cefdinir or any other part of this drug. **TELL YOUR DOCTOR:** If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. This drug may interact with other drugs or health problems. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well. Take with or without food. Take with food if it causes an upset stomach. Do not take antacids that have aluminum or magnesium in them within 2 hours of this drug. Do not take iron products within 2 hours before or 2 hours after taking this drug. Shake well before use. Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug. You may give this drug with infant formula that has iron in it. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store liquid (suspension) at room temperature. Throw away any part not used after 10 days. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Do not use longer than you have been told. A second infection may happen. If you have high blood sugar (diabetes) and test your urine glucose, talk with your doctor to find out which tests are best to use. If you have high blood sugar (diabetes), talk with the doctor. Some of these products have sugar. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. If you take this drug with an iron product, your stools may turn a reddish color. This is normal. Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY?
WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Not able to pass urine or change in how much urine is passed. Dark urine or yellow skin or eyes. Seizures. Fever, chills, or sore throat; any unexplained bruising or bleeding; or feeling very tired or weak. Diarrhea is common with antibiotics. Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Diarrhea. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

Copyright 2019 CDI, LLC. All rights reserved.
All rights reserved.
Issue Date: December 11, 2019

This information should not be used to decide whether or not to take this medicine or any other medicine. Only your health care provider has the